

On knots, braids and Gambaudo-Ghys quasi-morphisms

Michael Brandenbursky

October 22, 2009

Abstract

We study quasi-morphisms on the group of compactly supported area-preserving diffeomorphisms of a closed two-dimensional disc. In this paper we discuss possible variations of Gambaudo-Ghys construction, using quasi-morphisms on the groups P_n of pure braids on n strings. There arises a natural question: How to build quasi-morphisms of geometric origin on P_n ? We show that for this purpose it is possible to use knot invariants which satisfy specific properties. In particular we study quasi-morphisms which come from knot Floer homology and Khovanov-type homology. We also compute values of the homogeneous quasi-morphisms, obtained by this construction, on the time-one flow of a generic time-independent Hamiltonian in terms of its Reeb graph.

1 Introduction

Real-valued quasi-morphisms are known to be a helpful tool in the study of algebraic structure of non-Abelian groups, especially the ones that admit a few or no (linearly independent) real-valued homomorphisms. Recall that a *quasi-morphism* on a group G is a function $\varphi : G \rightarrow \mathbb{R}$ which satisfies the homomorphism equation up to a bounded error: there exists $K_\varphi > 0$ such that

$$|\varphi(ab) - \varphi(a) - \varphi(b)| \leq K_\varphi$$

for all $a, b \in G$. A quasi-morphism φ is called *homogeneous* if $\varphi(a^m) = m\varphi(a)$ for all $a \in G$ and $m \in \mathbb{Z}$. Any quasi-morphism φ can be *homogenized*: setting

$$\tilde{\varphi}(a) := \lim_{k \rightarrow +\infty} \varphi(a^k)/k \tag{1}$$

we get a homogeneous (possibly trivial) quasi-morphism $\tilde{\varphi}$.

The group $\mathcal{D}$ of compactly supported area-preserving diffeomorphisms of a closed two-dimensional disc is known to admit many (linearly independent) homogeneous quasi-morphisms (see e.g. [4], [12], [6]). In this paper we consider a particular construction, due to Gambaudo and Ghys [12], which produces quasi-morphisms on $\mathcal{D}$ from quasi-morphisms on the groups P_n of pure braids on n strings. The quasi-morphisms on P_n are constructed, in turn, from certain invariants of n -component links in $\mathbb{R}^3$ in the following way: close up a pure braid in a link in the standard way and take the value of the invariant on that link. The link invariant used by Gambaudo and Ghys is the signature. To show that the homogenization of the resulting quasi-morphism on $\mathcal{D}$ is non-trivial, Gambaudo and Ghys evaluate it on a diffeomorphism generated by a (compactly supported) time-independent radially symmetric Hamiltonian function on the disc (i.e. on a diffeomorphism which is the time-1 map of the Hamiltonian flow generated by the Hamiltonian function).

In this paper we discuss possible variations of the Gambaudo-Ghys construction involving other link and knot invariants. In particular, we formulate sufficient conditions for a knot invariant to yield a quasi-morphism on the full braid group B_n on n strings and so on P_n . We also consider two specific remarkable knot/link invariants: the Rasmussen link invariant s [25], which comes from a Khovanov-type theory, and the knot invariant τ [23], which comes from the Ozsvath-Szabo knot Floer homology. In [2] Baader has shown that the Rasmussen invariant s defines a quasi-morphism on B_n . We show that the homogenization of this quasi-morphism is equal to the classical linking number homomorphism on B_n . We also show that the situation with the Ozsvath-Szabo invariant τ is similar: it defines a quasi-morphism on B_n and its homogenization is again the linking number homomorphism multiplied by 2. In both cases the resulting homogeneous quasi-morphisms on $\mathcal{D}$ are homomorphisms, equal, up to a non-zero multiplicative constant, to the Calabi homomorphism.

We also discuss the computation of the quasi-morphisms on $\mathcal{D}$, obtained by Gambaudo-Ghys construction, on a diffeomorphism generated by a **time-independent** (compactly supported) Hamiltonian H . For a generic Hamiltonian H of this sort we present the result of the computation in terms of the Reeb graph of H and the integral of the push-forward of H to the graph against a certain signed measure on the graph. This result enables us to show that the Calabi homomorphism and the induced signature quasi-morphism on $\mathcal{D}$ are asymptotically equivalent on flows generated by time-independent (compactly supported) Hamiltonians.

The problem of computing any of the Gambaudo-Ghys quasi-morphisms on other elements of $\mathcal{D}$ seems to be much more difficult (see Section 6 for a related question).

Plan of the paper. In Section 2 we discuss the Gambaudo-Ghys construction. In Section 3 we formulate sufficient conditions for a knot invariant to yield a quasi-morphism on the full braid group B_n . In Section 4 we give examples of such knot invariants and homogeneous quasi-morphisms on the group $\mathcal{D}$ defined by them. In

Section 5 we define a set of generic time-independent Hamiltonians, and discuss the computation of the quasi-morphisms (defined by knot invariants) on the elements of $\mathcal{D}$ generated by these Hamiltonians. In Section 6 we discuss some open problems.

Acknowledgments. First of all, I would like to thank Michael Entov, who has introduced me to this subject, guided and helped me a lot while I was working on this paper. I would like to thank Michael Polyak for his valuable suggestions in the course of my work on this paper. I would like to thank Leonid Polterovich for helpful comments and for giving me the opportunity to present a part of this work at the Geometry and Dynamics seminar in Tel Aviv University.

2 Gambaudo-Ghys construction

In this section we recall the construction of homogeneous quasi-morphisms on the group $\mathcal{D} = \text{Diff}^\infty(\mathbb{D}^2, \partial\mathbb{D}^2, \text{area})$ of area-preserving diffeomorphisms of the closed unit disc $\mathbb{D}^2 \subset \mathbb{R}^2$ identical near the boundary. Denote the space of Hamiltonians $H : \mathbb{D}^2 \times [0, 1] \rightarrow \mathbb{R}$, such that the support of $H_t := H(\cdot, t)$ is compact for any t , by $\mathcal{H}$. We will denote the space of autonomous (i.e. time-independent) Hamiltonians by $\mathcal{H}_0$.

Let us now recall a few preliminaries concerning quasi-morphisms and braids.

The *defect* of a quasi-morphism φ on a group G is a minimal positive number D_φ , such that $|\varphi(gh) - \varphi(g) - \varphi(h)| \leq D_\varphi$ for all $g, h \in G$.

Throughout the paper the induced homogeneous quasi-morphism obtained by the homogenization of φ (see (1)) will be denoted by $\tilde{\varphi}$.

Denote by B_n the group of braids on n strands. The pure braid group $P_n \subset B_n$ can be identified with the fundamental group $\pi_1(X_n, \bar{z})$ of the space X_n of the ordered n -tuples of distinct points in $\mathbb{D}^2$, where $\bar{z} = (z_1, \dots, z_n) \in X_n$ is a base point in X_n .

Denote by $\sigma_1, \dots, \sigma_{n-1}$ the standard generators of B_n . The *braid length* $l(\gamma)$ of $\gamma \in B_n$ is the length of the shortest word representing γ with respect to the generators $\sigma_1, \dots, \sigma_{n-1}$.

We will work with homogeneous quasi-morphisms on P_n which are restrictions of homogeneous quasi-morphisms on B_n , or, equivalently (see e.g. [15]), with homogeneous quasi-morphisms $\tilde{\varphi} : P_n \rightarrow \mathbb{R}$ such that $\tilde{\varphi}(h) = \tilde{\varphi}(g^{-1}hg)$ for all $g \in B_n$ and $h \in P_n$. Such a quasi-morphism on P_n will be called *B_n -invariant*. This is a technical condition—the results below are likely to hold for all homogeneous quasi-morphisms on P_n .

The following construction, due to Ghys and Gambaudo [12], takes a B_n -invariant homogeneous quasi-morphism on P_n and produces from it a quasi-morphism on $\mathcal{D}$.

Let $\bar{x} = (x_1, \dots, x_n)$ be any point in X_n . Take $g \in \mathcal{D}$ and any path g_t , $0 \leq t \leq 1$, in $\mathcal{D}$ between Id and g . Connect $\bar{z}$ to $\bar{x}$ by the straight line in $(\mathbb{D}^2)^n$, then act on $\bar{x}$ with the path g_t , and then connect $g(\bar{x})$ to $\bar{x}$ by the straight line in $(\mathbb{D}^2)^n$. We get a loop in $(\mathbb{D}^2)^n$.

In coordinates it looks as follows. Connect z_i to x_i by straight lines $\mathfrak{l}_{1,i} : [0, \frac{1}{3}] \rightarrow \mathbb{D}^2$ in the disc, then act with the path $g_{3t-1} : [\frac{1}{3}, \frac{2}{3}] \rightarrow \mathbb{D}^2$, on each x_i , and then connect $g(x_i)$ to z_i by straight lines $\mathfrak{l}_{2,i} : [\frac{2}{3}, 1] \rightarrow \mathbb{D}^2$ in the disc, for all $1 \leq i \leq n$. It is a well known fact that $\mathcal{D}$ coincides with the group of compactly supported Hamiltonian diffeomorphisms of $\mathbb{D}^2$ (see e.g. [20]). Thus any identity-based path $\{h_t\}$ in $\mathcal{D}$, $0 \leq t \leq 1$, can be viewed as a Hamiltonian flow generated by a (time-dependent) Hamiltonian $H : \mathbb{D}^2 \times [0, 1] \rightarrow \mathbb{R}$, such that the support of $H_t := H(\cdot, t)$ is compact for any t . It follows that for almost all n -tuples of different points $x_1, \dots, x_n$ in the disc the concatenations of the paths $\mathfrak{l}_{1,i} : [0, \frac{1}{3}] \rightarrow \mathbb{D}^2$, $g_{3t-1} : [\frac{1}{3}, \frac{2}{3}] \rightarrow \mathbb{D}^2$ and $\mathfrak{l}_{2,i} : [\frac{2}{3}, 1] \rightarrow \mathbb{D}^2$, $i = 1, \dots, n$, yield a loop in X_n . The homotopy type of this loop is an element in P_n , which is independent of the choice of g_t because $\mathcal{D}$ is contractible [10]. We will denote this element by $\gamma(g; \bar{x})$.

Let $\tilde{\varphi}$ be a B_n -invariant homogeneous quasi-morphism on P_n . Set

$$\Phi(g) = \int_{X_n} \tilde{\varphi}(\gamma(g; \bar{x})) d\bar{x}, \quad (2)$$

where $d\bar{x} = dx_1 \cdot \dots \cdot dx_n$.

Lemma 2.1 (c.f. [12]). *The function Φ is well defined. It is a quasi-morphism on $\mathcal{D}$.*

As an immediate corollary we get that the formula

$$\tilde{\Phi}(g) := \lim_{p \rightarrow \infty} \frac{1}{p} \int_{X_n} \tilde{\varphi}(\gamma(g^p; \bar{x})) d\bar{x}$$

yields a homogeneous quasi-morphism $\tilde{\Phi} : \mathcal{D} \rightarrow \mathbb{R}$.

Proof. Step 1. Let us prove that $|\Phi(h)| < \infty$ for $h \in \mathcal{D}$.

For any isotopy h_t , $0 \leq t \leq 1$, in $\mathcal{D}$ between $h = h_1$ and Id , and any $\bar{x} \in X_n$ denote

$$l_{i,j} := \frac{h_t(x_i) - h_t(x_j)}{\|h_t(x_i) - h_t(x_j)\|} : [0, 1] \rightarrow \mathbb{S}^1.$$

Denote by

$$L_{i,j}(\bar{x}) = \frac{1}{2\pi} \int_0^1 \left\| \frac{\partial}{\partial t} (l_{i,j}) \right\| dt, \quad (3)$$

where $\|\cdot\|$ is the Euclidean norm. Note that $L_{i,j}(\bar{x})$ is the length of the path $l_{i,j}$ divided by 2π . Thus we get the following inequality

$$\sum_{i < j}^n 2(L_{i,j}(\bar{x}) + 4) \geq l(\gamma(h; \bar{x})). \quad (4)$$

Denote the extension of the B_n -invariant homogeneous quasi-morphism $\tilde{\varphi} : P_n \rightarrow \mathbb{R}$ to a homogeneous quasi-morphism on B_n by the same symbol $\tilde{\varphi}$.

Note that for any homogeneous quasi-morphism $\tilde{\varphi} : B_n \rightarrow \mathbb{R}$ one has

$$|\tilde{\varphi}(\gamma)| \leq \left(D_{\tilde{\varphi}} + \max_{1 \leq i \leq n-1} |\tilde{\varphi}(\sigma_i)| \right) l(\gamma). \quad (5)$$

It follows from (5) that:

$$l(\gamma(h; \bar{x})) \geq \frac{|\tilde{\varphi}(\gamma(h; \bar{x}))|}{N}, \quad (6)$$

where $N = D_{\tilde{\varphi}} + \max_{1 \leq i \leq n-1} |\tilde{\varphi}(\sigma_i)|$.

Inequalities (4) and (6) yield the following inequality:

$$\sum_{i < j}^n 2(L_{i,j}(\bar{x}) + 4) \geq \frac{|\tilde{\varphi}(\gamma(h; \bar{x}))|}{N}. \quad (7)$$

It follows that

$$\begin{aligned} |\Phi(h_1)| &\leq 2N \cdot \text{vol}((\mathbb{D}^2)^{n-2}) \sum_{i < j}^n \int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{2\pi} \int_0^1 \left\| \frac{\partial}{\partial t} (l_{i,j}) \right\| dt dx_i dx_j + \\ &\quad + 4Nn(n+1) \text{vol}((\mathbb{D}^2)^n). \end{aligned}$$

Denote by $h_t^{i,j} = h_t(x_i) - h_t(x_j)$ and note that

$$\left\| \frac{\partial}{\partial t} (l_{i,j}) \right\| = \left\| \frac{\frac{\partial}{\partial t} h_t^{i,j}}{\|h_t^{i,j}\|} - \frac{h_t^{i,j} \cdot \frac{\partial}{\partial t} h_t^{i,j}}{\|h_t^{i,j}\|^3} \right\|.$$

It follows from Cauchy-Schwartz inequality that

$$\left\| \frac{\partial}{\partial t} (l_{i,j}) \right\| \leq \frac{2 \left\| \frac{\partial}{\partial t} (h_t^{i,j}) \right\|}{\|h_t^{i,j}\|}.$$

Let $H \in \mathcal{H}$ be a Hamiltonian generating the flow h_t , $0 \leq t \leq 1$.

Note that $\left\| \frac{\partial h_t(x)}{\partial t} \right\| = \|(\nabla H_t)(h_t(x))\| \leq \max_{x \in \mathbb{D}^2, t \in [0,1]} \|(\nabla H_t)(x)\|$. Denote by

$$M_H = \max_{x \in \mathbb{D}^2, t \in [0,1]} \|(\nabla H_t)(x)\|.$$

Thus:

$$\left\| \frac{\partial}{\partial t} (l_{i,j}) \right\| \leq \frac{4M_H}{\|h_t(x_i) - h_t(x_j)\|}.$$

Recall that h_t is a Hamiltonian flow and so is area preserving. Then:

$$\int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{2\pi} \int_0^1 \left\| \frac{\partial}{\partial t} (l_{i,j}) \right\| dt dx_i dx_j \leq \frac{4M_H}{2\pi} \left(\int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{\|x_i - x_j\|} dx_i dx_j \right).$$

Note that $\int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{\|x_i - x_j\|} dx_i dx_j$ is well defined, because by Fubini-Tonelli theorem

$$\int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{\|x_i - x_j\|} dx_i dx_j \leq \int_{\mathbb{D}^2} \left(\int_{B_\delta(x_j)} \frac{1}{\|x_i - x_j\|} dx_i + \int_{\mathbb{D}^2 \setminus (\mathbb{D}^2 \cap B_\delta(x_j))} \frac{1}{\|x_i - x_j\|} dx_i \right) dx_j,$$

where $\delta \leq \frac{1}{2}$ is some positive constant. Thus by using the polar coordinates on $B_\delta(x_j)$ we get

$$\int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{\|x_i - x_j\|} dx_i dx_j < \int_{\mathbb{D}^2} \left(2\pi\delta + \frac{\text{vol}(\mathbb{D}^2)}{\delta} \right) dx_j = \pi^2 \left(2\delta + \frac{1}{\delta} \right).$$

Denote by $N' = 4Nn(n+1) \left(\frac{\text{vol}((\mathbb{D}^2)^{n-2})}{2\pi} \left(\int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{\|x_i - x_j\|} dx_i dx_j \right) + \text{vol}((\mathbb{D}^2)^n) \right)$, then

$$|\Phi(h)| \leq N' \cdot M_H.$$

Step 2. Let us prove that Φ is a quasi-morphism. Let g_1 and g_2 be elements of $\mathcal{D}$. Note that for almost all $\bar{x} \in X_n$

$$\gamma(g_1 g_2; \bar{x}) = \gamma(g_1; g_2(\bar{x})) \cdot \gamma(g_2; \bar{x})$$

The diffeomorphisms g_1 and g_2 are area-preserving. Thus

$$|\Phi(g_1 g_2) - \Phi(g_1) - \Phi(g_2)| = \left| \int_{X_n} \tilde{\varphi}_n(\gamma(g_1 g_2; \bar{x})) - \tilde{\varphi}_n(\gamma(g_1; g_2(\bar{x}))) - \tilde{\varphi}_n(\gamma(g_2; \bar{x})) d\bar{x} \right|.$$

It follows that

$$|\Phi(g_1 g_2) - \Phi(g_1) - \Phi(g_2)| \leq \int_{X_n} D_{\tilde{\varphi}}.$$

□

From now on if we have a B_n -invariant homogeneous quasi-morphisms $\tilde{\varphi}_n : P_n \rightarrow \mathbb{R}$, then the induced homogeneous quasi-morphisms on $\mathcal{D}$ will be denoted by $\tilde{\Phi}_n$.

3 Quasi-morphisms on B_n

Given a braid $\gamma \in B_n$, we denote by $\widehat{\gamma}$ the link obtained by closing up the braid in the standard way. See Figure 1 for an example.

One of the ways to find a quasi-morphism on B_n is to find an $\mathbb{R}$ -valued invariant I of (isotopy classes of) links in S^3 so that

$$|I(\widehat{\alpha\beta}) - I(\widehat{\alpha}) - I(\widehat{\beta})| \leq K_I,$$

where $K_I > 0$ depends only on I . An example of such an invariant is the link signature – this was proved in [12].

In this section we will formulate a sufficient condition for a knot invariant under which it yields a quasi-morphism on B_n . As a corollary we will define another quasi-morphism on B_n using the knot invariant τ , which comes from the knot Floer homology. First we will recall some notations from the knot theory.

The four-ball genus of a knot K in S^3 is the minimal genus of an oriented surface with boundary, which is embedded in $\mathbb{D}^4$ and whose boundary is $K \subset S^3 = \partial\mathbb{D}^4$. It will be denoted by $g_4(K)$. We denote by $\text{Conc}(S^3)$ the concordance group of knots in S^3 . For any knot K , K^* denotes its mirror image, and $-K$ denotes the same knot with the reversed orientation. For any two knots K_1 and K_2 the knot $K_1 \# K_2$ represents their connected sum.

Definition 3.1. A braid $\alpha \in B_n$ is called **positive** if it can be written as a product of only non-negative powers of the standard generators of B_n .

For each $k \in \mathbb{N}$ we will denote by L_k the set of all words written as a product of only non-negative powers of standard generators of B_n of the length k . We will define a total order on L_1 by setting

$$\sigma_1 \prec \sigma_2 \prec \dots \prec \sigma_{n-1}.$$




Figure 1: On the left is presented the braid σ_1^2 and on the right is presented the positive Hopf link $\widehat{\sigma_1^2}$.

This order induces a total lexicographic order $\prec$ on L_k for any $k \in \mathbb{N}$. Thus in each subset of L_k there exist a unique word which is minimal with respect to $\prec$.

For any braid $\beta \in B_n$ there exists a unique positive braid $\alpha_\beta \in B_n$ which satisfies the following properties:

1. The closure of $\widehat{\alpha_\beta \beta}$ is a knot.
2. The braid α_β is represented by a unique minimal word in the set L_k , where

$$k = \min_{\alpha \in B_n} \{l(\alpha) \mid \widehat{\alpha \beta} \text{ is a knot, } \alpha \text{ is a positive braid}\}.$$

Definition 3.2. Given a knot invariant I , we define a function

$$I : B_n \rightarrow \mathbb{R},$$

by $I(\beta) = I(\widehat{\alpha_\beta \beta})$, where α_β is defined above.

For example $I|_{P_n}$ is defined as follows: $I(\beta) = I(\widehat{\alpha \beta})$, where $\alpha = \sigma_1 \cdot \dots \cdot \sigma_{n-1}$.

Theorem 3.3. Suppose that a knot invariant I defines a homomorphism

$$I : \text{Conc}(S^3) \rightarrow \mathbb{R},$$

such that $|I(K)| \leq cg_4(K)$, where c is some positive real constant independent of K . Then I defines a quasi-morphism on P_n .

Proof. To simplify the notation we will prove here that $I|_{P_n}$ is a quasi-morphism. In the same way it is easy to prove that $I : B_n \rightarrow \mathbb{R}$ is a quasi-morphism.

Take any $\beta, \gamma \in P_n$. Note that there is a saddle cobordism S_1 with $\chi(S_1) = -1$ between $(\widehat{\alpha \gamma \# \alpha \beta})$ and $(\widehat{\alpha \gamma} \sqcup \widehat{\alpha \beta})$. It is easy to see that $\widehat{\alpha}$ is an unknot, so there is a cobordism S_2 between $\widehat{\alpha}$ and a closure of an identity braid e with $\chi(S_2) = 1 - n$. Then there is cobordism S_3 between $(\widehat{\gamma} \sqcup \widehat{\alpha})$ and $\widehat{\gamma} \sqcup \widehat{e}$, such that $\chi(S_3) = 1 - n$. See Figure 2.

In the proof of Theorem 2 in [2] Baader showed that for any 2 braids μ and ν in B_n there is a cobordism of an Euler characteristic $-n$ between $\widehat{\mu \nu}$ and $(\widehat{\mu} \sqcup \widehat{\nu})$. Thus there is a cobordism S_4 between $\widehat{\alpha \gamma}$ and $\widehat{\gamma}$, such that $\chi(S_4) = 1 - 3n$. And so there is a cobordism S_5 between $(\widehat{\alpha \gamma} \sqcup \widehat{\alpha \beta})$ and $\widehat{\alpha \beta \gamma}$, such that $\chi(S_5) = 1 - 4n$. So $S = S_5 S_1$ is a cobordism between $(\widehat{\alpha \gamma \# \alpha \beta})$ and $\widehat{\alpha \beta \gamma}$, such that $\chi(S) = -4n$. It means that the induced cobordism $\widehat{S}$ between $(-\widehat{\alpha \beta \gamma}^*) \# (\widehat{\alpha \gamma \# \alpha \beta})$ and $(-\widehat{\alpha \beta \gamma}^*) \# \widehat{\alpha \beta \gamma}$ has an Euler characteristic $\chi(\widehat{S}) = -4n$.

A knot K in $S^3 = \partial \mathbb{D}^4$ is called *slice* if there exists an embedding of the standard disc $\mathbb{D}^2 \subset \mathbb{R}^2$ in $\mathbb{D}^4$ such that the image of an embedding of $\partial \mathbb{D}^2$ is the knot K . It is a well known fact that for any knot K the knot $K \# -K^*$ is a slice knot. Thus

$$(-\widehat{\alpha \beta \gamma}^*) \# \widehat{\alpha \beta \gamma}$$




Figure 2: Cobordism S_3 between $-(\hat{\gamma}^* \sqcup \hat{\alpha}^*)$ and $\hat{\gamma}^* \sqcup \hat{e}^*$.

is a slice knot. Therefore

$$g_4(-(\widehat{\alpha\beta\gamma})^* \# (\widehat{\alpha\gamma} \# \widehat{\alpha\beta})) \leq 2n,$$

where g_4 is the four ball genus. We know that for every knot K the following inequality $|I(K)| \leq cg_4(K)$. This yields:

$$|I(-(\widehat{\alpha\beta\gamma})^* \# (\widehat{\alpha\gamma} \# \widehat{\alpha\beta}))| \leq c \cdot 2n.$$

Recalling that $I(-K^*) = -I(K)$ and $I(K_1 \# K_2) = I(K_1) + I(K_2)$ finishes the proof. $\square$

Remark 3.4. Note that g_4 defines a norm on $\text{Conc}(S^3)$. Theorem 3.3 can be reformulated like this. Each element of $\text{Hom}(\text{Conc}(S^3), \mathbb{R})$, which is Lipschitz with respect to this norm defines a quasi-morphism on B_n .

In the next section we will present the following three knot/link invariants: the Ozsvath-Szabo knot invariant τ , the Rasmussen link invariant s , and the link signature. It is well known that all of them satisfy the conditions of Theorem 3.3. Thus by Theorem 3.3 they define quasi-morphisms on B_n .

4 Examples

4.1 Signature quasi-morphism

Now we will recall the definition of the signature link invariant. Let L be an oriented link in S^3 , then by the Seifert algorithm [18] there exists an oriented surface S_L with boundary L . Choose a basis $\{a_1, \dots, a_{2g+|L|-1}\}$ in $H_1(S_L, \mathbb{Z})$, then there exists a symmetric bilinear form on $H_1(S_L, \mathbb{Z})$, which is defined as follows:

$$\Omega(a_i, a_j) = lk(a_i, a_j^+) + lk(a_j, a_i^+),$$

where lk is the linking number and a_i^+ is a push of the curve, which represents a_i in S_L , from S_L along the positive normal direction to S_L . Tensoring by $\mathbb{R}$ we get a symmetric bilinear form on $H_1(S_L, \mathbb{R})$. The signature of this form is independent of the choices of the Seifert surface and the basis of $H_1(S_L, \mathbb{Z})$, see [18]. Thus it is an invariant of L and is denoted by $sign(L)$. The following proposition was proved by Gambaudo and Ghys in [12]. Our Theorem 3.3 gives another proof of this proposition.

Proposition 4.1. *Link signature defines a quasi-morphism on B_n .*

We will denote the induced homogeneous quasi-morphism on B_n by $\widetilde{sign}_n$. The induced homogeneous quasi-morphism on $\mathcal{D}$ is denoted by $\widetilde{Sign}_{n, \mathbb{D}^2}$.

Proposition 4.2. 1. *For every $n > 2$ the homogeneous quasi-morphism $\widetilde{sign}_n : B_n \rightarrow \mathbb{R}$ is not a homomorphism.*

2. *The restriction of $\widetilde{sign}_n$ to P_n is not a homomorphism.*

The proof will be given in Section 5.

Theorem 4.3. *For each $g \in \mathcal{D}$ generated by an autonomous Hamiltonian*

$$\lim_{n \rightarrow \infty} \frac{\widetilde{Sign}_{n, \mathbb{D}^2}(g)}{\pi^{n-1} n(n-1)} = -\mathcal{C}(g).$$

The proof will be given in Section 5.

4.2 Rasmussen quasi-morphism

The Rasmussen invariant $s(K)$ of a knot K in S^3 was discovered by Jacob Rasmussen in 2004 (see [25]). This is a very powerful knot invariant, which, for example, gives a lower bound on the four-ball genus of a knot. It comes from the Lee theory, which is closely related to the Khovanov homology. For more information see [16] and [17]. Recently this invariant was extended to links by Beliakova and Wehrli [5]. It has many interesting properties. We list here eight of them:

$$s(K) = s(-K), \tag{8}$$

$$s(K_1 \# K_2) = s(K_1) + s(K_2), \tag{9}$$

$$s(K^*) = -s(K), \tag{10}$$

$$|s(K)| \leq 2g_4(K), \tag{11}$$




Figure 3: On the left there is a positive crossing and on the right there is a negative crossing.

$$s(K_-) \leq s(K_+) \leq s(K_-) + 2, \quad (12)$$

where K_+ and K_- are knots, which differ by a single crossing and this crossing is positive in K_+ and negative in K_- (see Figure 3).

For a positive link L

$$s(L) = w(D(L)) - o(D(L)) + 1, \quad (13)$$

where $D(L)$ is some positive diagram of L , $w(D(L))$ is the number of positive crossings in $(D(L))$ minus the number of negative crossings in $(D(L))$, and $o(D(L))$ is the number of Seifert circles, when the Seifert algorithm is applied to $D(L)$.

$$1 + w(D(K)) - o(D(K)) \leq s(K) \leq -1 + w(D(K)) + o(D(K)), \quad (14)$$

where $D(K)$ is some diagram of a knot K (see [2]). In fact, it is shown in [2] that property (14) follows from properties (10), (12) and (13). For an alternating knot K

$$s(K) = -\text{sign}(K), \quad (15)$$

where sign is the signature knot invariant.

In his paper [2] Baader proved that the Rasmussen link invariant s defines a quasi-morphism on the braid group B_n , with a defect bounded by $n + 1$. Alternatively, we see that s satisfies properties (8), (9), (10) and (11). Thus our Theorem 3.3 proves that s defines a quasi-morphism on B_n . The induced family of homogeneous quasi-morphisms on $\mathcal{D}$ is well defined and is denoted by $\widetilde{Ras}_n$.

Theorem 4.4. *Suppose that $\varphi : B_n \rightarrow \mathbb{R}$ is a quasi-morphism defined by a link invariant, which satisfies property (14). Then $\widetilde{\varphi} = lk$.*




Figure 4: The braid $\beta = \sigma_1 \sigma_2 \dots \sigma_{n-1}$ and its closure is an unknot.

Proof. Take $\alpha \in B_n$. Then for all $p > 0$ there exists a positive braid β_p in B_n , such that $\widehat{\alpha^p \beta_p}$ is a knot and $l(\beta_p) \leq L(n)$, where l is a braid length and $L(n)$ depends only on n . For example if $\alpha \in P_n$, then we can take $\beta_p = \beta = \sigma_1 \cdot \dots \cdot \sigma_{n-1}$ for all $p \in \mathbb{N}$. See Figure 4. Note that $w(D(\widehat{\alpha^p \beta_p})) = lk(\alpha^p \beta_p)$. It follows from the property (14) that:

$$1 + lk(\alpha^p \beta_p) - n \leq \varphi(\alpha^p \beta_p) \leq -1 + lk(\alpha^p \beta_p) + n.$$

For all $p > 0$ there exist a constant $K(n) > 0$, such that $|\varphi(\beta_p)| < K(n)$. Also note that the following inequality $|lk(\beta_p)| \leq L(n)$ is true for all $p > 0$, where $L(n)$ is some positive real constant. Thus:

$$\lim_{p \rightarrow \infty} \frac{1}{p} (1 + lk(\alpha^p \beta_p) - n) \leq \lim_{p \rightarrow \infty} \frac{1}{p} \varphi(\alpha^p \beta_p) \leq \lim_{p \rightarrow \infty} \frac{1}{p} (-1 + lk(\alpha^p \beta_p) + n).$$

Therefore $\lim_{p \rightarrow \infty} \frac{1}{p} \varphi(\alpha^p \beta_p) = lk(\alpha)$. Now φ is a quasi-morphism with a defect K_φ and so

$$|\varphi(\alpha^p \beta_p) - \varphi(\alpha^p)| \leq |\varphi(\beta_p)| + K_\varphi < K(n) + K_\varphi.$$

It follows that

$$lk(\alpha) = \lim_{p \rightarrow \infty} \frac{\varphi(\alpha^p \beta_p)}{p} = \lim_{p \rightarrow \infty} \frac{\varphi(\alpha^p)}{p} = \tilde{\varphi}(\alpha).$$

□

Definition 4.5. A link diagram is called **alternating** if the crossings alternate under, over, under, over, and so on as one travels along each component of the link. A link is called **alternating** if it has an alternating diagram. A braid $\alpha \in B_n$ is called **alternating** if $\widehat{\alpha}$ is an alternating link.

Corollary 4.6. Each $\alpha \in B_n$ satisfies $\tilde{s}(\alpha) = lk(\alpha)$. If $\gamma \in B_n$ is an alternating braid, then $\widetilde{sign}(\gamma) = -lk(\gamma)$.

Proof. The first statement follows directly from Theorem 4.4.

Now we will prove the second statement. For each $p \in \mathbb{N}$ the braid γ^p is alternating and there exists an alternating braid β_p such that $l(\beta_p) < n$ and $\widehat{\gamma^p \beta_p}$ is an alternating knot. It follows from property (15) that

$$\lim_{p \rightarrow \infty} \frac{\text{sign}(\gamma^p \beta_p)}{p} = - \lim_{p \rightarrow \infty} \frac{\text{lk}(\gamma^p \beta_p)}{p} = -\text{lk}(\gamma).$$

Using the fact that sign is a quasi-morphism we get the following equality

$$\lim_{p \rightarrow \infty} \frac{\text{sign}(\gamma^p \beta_p)}{p} = \lim_{p \rightarrow \infty} \frac{\text{sign}(\gamma^p)}{p} = \widetilde{\text{sign}}(\gamma).$$

□

In [2] Baader defined the following quasi-morphism:

$$s - \text{lk} + n - 1 : B_n \rightarrow \mathbb{R}.$$

This quasi-morphism maps all positive braids to zero. Therefore it descends to the quasi-morphism on $B_n / \langle \Delta_n \rangle$, where Δ_n^2 is the positive generator of the center $Z(B_n)$ of B_n . In [2] Baader asked whether this quasi-morphism is bounded for $n > 2$. Here we give a positive answer to this question.

Proposition 4.7. *The quasi-morphism $s - \text{lk} + n - 1 : B_n \rightarrow \mathbb{R}$ is bounded.*

Proof. Take any braid $\beta \in B_n$ and the positive braid α_β defined in Section 3. Using the fact that s is a quasi-morphism, $s(\alpha_\beta) \leq n - 1$, and $\text{lk}(\alpha_\beta) \leq n - 1$ we get the following inequality

$$|s(\beta) - \text{lk}(\beta)| \leq |s(\beta \alpha_\beta) - \text{lk}(\beta \alpha_\beta)| + 3(n - 1).$$

It follows from property (14) that

$$|s(\beta \alpha_\beta) - \text{lk}(\beta \alpha_\beta)| \leq n - 1.$$

This yields:

$$|s(\beta) - \text{lk}(\beta) + n - 1| \leq 5(n - 1).$$

The last inequality shows that this quasi-morphism is bounded on B_n and thus also on $B_n / \langle \Delta_n \rangle$. □

The proof of the following theorem is almost identical to the proof of Theorem 2.4 in [11] by Gambaudo and Ghys. It will be presented in Section 5.

Theorem 4.8. *For every $g \in \mathcal{D}$*

$$\widetilde{\text{Ras}}_n(g) = 2n(n - 1)\pi^{n-1} \cdot \mathcal{C}(g). \quad (16)$$

4.3 τ quasi-morphism and the braid number of a knot.

In [23] Ozsvath and Szabo defined a knot invariant τ coming from the knot Floer homology. This invariant satisfies in particular the following properties:

$$\tau(K^*) = -\tau(K), \quad (17)$$

$$\tau(K_1 \# K_2) = \tau(K_1) + \tau(K_2), \quad (18)$$

$$\tau(-K) = \tau(K), \quad (19)$$

$$|\tau(K)| \leq g_4(K), \quad (20)$$

$$0 \leq \tau(K^+) - \tau(K^-) \leq 1, \quad (21)$$

$$\tau(\widehat{\alpha}) = \frac{lk(\alpha) - n + 1}{2}, \quad (22)$$

where $\alpha \in B_n$ is a positive braid, such that $\widehat{\alpha}$ is a knot.

The first five properties were proved in the paper of Ozsvath and Szabo [23] and the last property was proved by Livingston in [19].

Proposition 4.9. *The knot invariant τ defines a quasi-morphism on B_n and*

$$2\widetilde{\tau} = lk.$$

Proof. The first statement follows directly from Theorem 3.3.

Note that property (14) follows from properties (17), (21) and (22), where s is substituted by 2τ . The lower bound is proved exactly as in [26] and the upper bound is true, because of the property (17) combined with the lower bound. Thus the second statement follows from the proof of Theorem 4.4. $\square$

Recall that every knot K in $\mathbb{R}^3$ can be presented as a closure of some braid in B_n . The braid number of K is the minimal such n . It is denoted by $br(K)$.

Theorem 4.10. *Let s and τ be the quasi-morphisms on B_n which are induced from Rasmussen and Ozsvath-Szabo knot invariants. Then for every $\alpha \in B_n$ the following inequality holds $|s(\alpha) - 2\tau(\alpha)| \leq 2(n - 1)$.*

Proof. As we explained before both s and 2τ satisfy property (14). It means that:

$$|s(\alpha) - lk(\beta_\alpha \alpha)| \leq n - 1 \quad \text{and} \quad |2\tau(\alpha) - lk(\beta_\alpha \alpha)| \leq n - 1.$$

Thus by triangle inequality $|s(\alpha) - 2\tau(\alpha)| \leq 2(n - 1)$. $\square$

Corollary 4.11. *For every knot K the following inequality holds:*

$$|s(K) - 2\tau(K)| \leq 2(br(K) - 1).$$

4.4 ω -signature quasi-morphisms

In this section we define for each $n \geq 2$ a family of quasi-morphisms on B_n and show that this family gives a basis for

$$\mathcal{S} = \{\tilde{\varphi} : A_n \rightarrow \mathbb{R} \mid \tilde{\varphi} \text{ is a homogeneous quasi-morphism on } B_n\},$$

where $A_n = \langle \eta_{2,n}^+, \eta_{2,n}^-, \dots, \eta_{n,n}^+, \eta_{n,n}^- \rangle$. Here $\eta_{i,n}^+$ is a pure braid in P_n , in which the i -th strand makes one twist in the positive direction around previous $i-1$ strands, and $\eta_{i,n}^-$ is a pure braid in P_n , in which the i -th strand makes one twist in the negative direction around previous $i-1$ strands.

As we will show in Section 5, in order to evaluate a homogeneous quasi-morphism on $\mathcal{D}$, defined by means of the Gambaudo-Ghys construction, on the time-one flow of a generic autonomous Hamiltonian, one only needs to know the values of the corresponding quasi-morphism on P_n on the set A_n .

Now we will remind the definition of the ω -signature link invariant, where $\omega \neq 1$ is a complex number. Let L be an oriented link in S^3 . Choose any Seifert surface S_L . Choose a basis $\{a_1, \dots, a_{2g+|L|-1}\}$ in $H_1(S_L, \mathbb{Z})$, then there exists a bilinear form B on $H_1(S_L, \mathbb{Z})$, which is defined as follows:

$$B(a_i, a_j) = lk(a_i, a_j^+),$$

where lk is the linking number and a_i^+ is a push of the curve, which represents a_i in S_L , from S_L along the positive normal direction to S_L . Tensoring this form by $\mathbb{C}$, we get a bilinear form on $H_1(S_L, \mathbb{C})$. The signature of the following hermitian form on $H_1(S_L, \mathbb{C})$

$$\Omega_\omega(a_i, a_j) = (1 - \omega)B(a_i, \overline{a_j}) + (1 - \overline{\omega})B(\overline{a_j}, a_i)$$

is independent of the choice of the Seifert surface and the base for $H_1(S_L, \mathbb{Z})$, see [18]. Thus for each ω this signature is a link invariant denoted by $sign_\omega(L)$.

Gambaudo and Ghys proved in [12] that $sign_\omega$ is a quasi-morphism on B_n . Alternatively, Theorem 3.3 shows that $sign_\omega$ is a quasi-morphism on B_n . The homogeneous quasi-morphism will be denoted as usual by $\widetilde{sign}_\omega$. Recall that the torus link $K(p, q)$ has a braid representation $(\sigma_1 \cdot \dots \cdot \sigma_{p-1})^q$.

Suppose that $\omega = e^{2\pi i \theta}$, where $\theta \in [0, 1] \cap \mathbb{Q}$. Then it follows from Proposition 5.1 and Proposition 5.2 in [13] that:

$$\widetilde{sign}_\omega(K(n, n)) = -2n(n - 2l + 1)\theta - 2l(l - 1) \quad (23)$$

for $\frac{l-1}{n} \leq \theta \leq \frac{l}{n}$, where $1 \leq l \leq n$.

Proposition 4.12. *Let $\omega = e^{2\pi i\theta}$, where $\theta \in [0, 1] \cap \mathbb{Q}$. Then:*

$$\widetilde{sign}_\omega(\eta_{i,n}) = \begin{cases} 4(1-i)\theta, & \text{if } 0 \leq \theta \leq \frac{1}{i}, \\ 4(1-l)(1-\theta), & \text{if } \frac{l-1}{i} \leq \theta \leq \frac{l-1}{i-1}, \quad 2 \leq l \leq i-1, \\ 4(l-i)\theta, & \text{if } \frac{l-1}{i-1} \leq \theta \leq \frac{l}{i}, \quad 2 \leq l \leq i-1, \\ 4(1-i)\theta, & \text{if } \frac{i-1}{i} \leq \theta \leq 1. \end{cases} \quad (24)$$

Proof. Note that $K(n, np) = \eta_{2,n}^p \cdot \dots \cdot \eta_{n,n}^p$. Thus

$$\widetilde{sign}_\omega(K(n, n)) = \widetilde{sign}_\omega(\eta_{2,n} \cdot \dots \cdot \eta_{n,n}) = \sum_{i=2}^n \widetilde{sign}_\omega(\eta_{i,n}).$$

Also note that for all $l, r \in \mathbb{N}$, such that $l \geq i$ and $r \geq i$, the following equation holds

$$\widetilde{sign}_\omega(\eta_{i,k}) = \widetilde{sign}_\omega(\eta_{i,r}).$$

It follows that:

$$\widetilde{sign}_\omega(\eta_{i,n}) = \widetilde{sign}_\omega(K(i, i)) - \widetilde{sign}_\omega(K(i-1, i-1)).$$

After a simple calculation combined with the last equality and (23) we get the result. $\square$

Proposition 4.13. *Denote by $\omega(\theta) = \omega^{2\pi i\theta}$. Then*

$$\mathcal{B} = \left\{ \widetilde{sign}_{\omega(\frac{1}{2})}, 2\widetilde{sign}_{\omega(\frac{1}{2})} - 3\widetilde{sign}_{\omega(\frac{1}{3})}, \dots, (n-1)\widetilde{sign}_{\omega(\frac{1}{n-1})} - n\widetilde{sign}_{\omega(\frac{1}{n})} \right\}$$

is a basis for $\mathcal{S}$.

Proof. If $n = 2$, it is obvious because $\widetilde{sign}_{\omega(\frac{1}{2})}(\eta_{2,2}) = -2$. Let $n > 2$. For $1 \leq i \leq n-1$, the equality: $\left((n-1)\widetilde{sign}_{\omega(\frac{1}{n-1})} - n\widetilde{sign}_{\omega(\frac{1}{n})} \right)(\eta_{i,n}) = 0$ follows from (24), and also we have the following equality:

$$\left((n-1)\widetilde{sign}_{\omega(\frac{1}{n-1})} - n\widetilde{sign}_{\omega(\frac{1}{n})} \right)(\eta_{n,n}) = 4.$$

Note also that $\widetilde{sign}_\omega(\eta_{i,i}) = \widetilde{sign}_\omega(\eta_{i,n})$. For $3 \leq i \leq n$ this yields

$$\left((i-1)\widetilde{sign}_{\omega(\frac{1}{i-1})} - i\widetilde{sign}_{\omega(\frac{1}{i})} \right)(\eta_{j,n}) = 0$$

for $2 \leq j \leq i - 1$ and

$$\left((i-1) \widetilde{\text{sign}}_{\omega(\frac{1}{i-1})} - i \widetilde{\text{sign}}_{\omega(\frac{1}{i})} \right) (\eta_{i,n}) = 4.$$

Thus $\mathcal{B}$ is a basis for $\mathcal{S}$.

□

5 Autonomous Hamiltonians

Definition 5.1. We say that a function $H \in \mathcal{H}_0$ is of **Morse type** if:

1. There exists a connected open neighborhood U of $\partial\mathbb{D}^2$, such that $\partial\overline{U} \setminus \partial\mathbb{D}^2$ is a smooth simple curve, $H|_{\overline{U}} \equiv 0$ and H has no degenerate critical points in $\mathbb{D}^2 \setminus \overline{U}$.
2. There exists a connected open neighborhood V of $\partial\mathbb{D}^2$, such that $V \supset \overline{U}$ and $H|_{V \setminus \overline{U}}$ has no critical points.

In this section we show that $\tilde{\Phi}$ is continuous in some sense on the space $\mathcal{H}_0$ (see Theorem 5.11).

It follows from Theorem 2.7 in [21] that Morse type Hamiltonians form a C^1 -dense subset of $\mathcal{H}_0$. For any Hamiltonian H in this subset we present a calculation of $\tilde{\Phi}$ on the time-one flow of H (Theorem 5.4). Our result is presented in terms of the integral of the push-forward of H to the Reeb graph of H against a certain signed measure on the graph.

Now let us explain how we associate to H and $\tilde{\varphi}$ the Reeb graph T_H of H and the signed measure $\mu_{\tilde{\varphi}}$ on T_H .

Definition 5.2. A **charged tree** T is a finite tree equipped with a signed measure, such that each edge has a total finite measure.

Given a tree T with l edges we will number them once and for all and denote them by $e_1, \dots, e_l$.

Now we are ready to explain how we associate to $(H, \tilde{\varphi})$ a charged tree $(T_H, \mu_{\tilde{\varphi}})$. Let $H \in \mathcal{H}_0$ be a Morse type function with r critical points in $\mathbb{D}^2 \setminus \overline{U}$, where U is as in Definition 5.1. Let $c = \partial\overline{U} \setminus \partial\mathbb{D}^2$ and suppose that the area of the domain $\overline{U}$ is $2\pi\varepsilon$ for some $\varepsilon > 0$.

Step 1. Here we recall the notion of angle-action symplectic coordinates.

Cut $\mathbb{D}^2$ along the singular level curves inside $\mathbb{D}^2 \setminus \overline{U}$, along the curve c , and along $\partial\mathbb{D}^2$. We get a_r open annuli A_i (a punctured disc is also viewed as an annulus) and an open annulus $A_0 = U^\circ$. Each A_i has two boundary components c_i and d_i , where $H|_{c_i} \leq H|_{d_i}$. By Liouville-Arnold theorem (see [1]) on each of the annuli A_i , there exist so-called angle-action symplectic coordinates (I_i, θ_i) , $\theta_i \in [0, 2\pi]$, and a C^∞ -function $h_i : [0, \frac{\text{area}(A_i)}{2\pi}] \rightarrow \mathbb{R}$, such that on each level curve δ_i in A_i one has $H|_{\delta_i} = h_i(I_i)$, where




Figure 5: Reeb graph which corresponds to the function H with 3 maximum points and 2 saddle points.

the coordinate I_i along δ_i is equal to the area of the annulus bounded by δ_i and c_i , divided by 2π . In these coordinates h_t moves points on each level curve with a constant speed $\frac{\partial \tilde{h}_i}{\partial I_i}$. Take the coordinates (I_0, θ_0) on A_0 , where $I_0 \in [\frac{1}{2} - \varepsilon, \frac{1}{2}]$ is the area coordinate divided by 2π , and θ_0 is the angle coordinate. Note that $\tilde{h}_0 = 0$, and

$$\int_{\mathbb{D}^2} H(p, q) dp dq = \sum_{i=1}^{a_r} \int_{A_i} \tilde{h}_i(I_i) dI_i d\theta_i = 2\pi \sum_{i=1}^{a_r} \int_{A_i} \tilde{h}_i(I_i) dI_i.$$

Step 2. Here we associate to the Morse type function $H \in \mathcal{H}_0$ its Reeb graph. Contract each connected component of each level set of H to a point. As a result one gets a graph called the Reeb graph of H . It will be denoted by T_H . One can check that in our case this is a tree. Its vertices correspond to the connected components of critical level sets of H . We will view T_H as a rooted tree with the root being the vertex of T_H corresponding to the domain $\bar{U}$.

Example 5.3. Figure 5 shows a Reeb graph with 5 edges and 6 vertices, such that 3 of them correspond to the maximum points, 2 of them correspond to the saddle points and the root corresponds to the region bounded by the curves c and $\partial \mathbb{D}^2$.

We call the Reeb graph *simple* if it has only 2 vertices and one edge. Note that the Reeb graph is simple if it corresponds to the monotone radial function, which is constant in the neighborhood of zero.

Step 3. The action coordinate on any of the annuli induces a coordinate I_j on the corresponding edge e_j of T_H . Thus each $H \in \mathcal{H}_0$ descends to a function $\tilde{h} : T_H \rightarrow \mathbb{R}$, where $\tilde{h}(x) = h_j(x)$, and $x \in e_j$, for each open edge e_j . Here $j \in \{0, \dots, a_r\}$. Let's define a signed measure $\mu_{\tilde{\varphi}}$ on T_H by setting

$$d\mu_{\tilde{\varphi}}(I_j) = (2\pi)^n \sum_{i=2}^n \tilde{\varphi}(\eta_{i,n}) i \binom{n}{i} (I_j)^{i-1} \left(\frac{1}{2} - I_j\right)^{n-i} dI_j,$$

for each $j \in \{0, \dots, a_r\}$, where I_j is a coordinate on each edge e_j .

Theorem 5.4. *Let $H \in \mathcal{H}_0$ be a Morse type function, and $\tilde{\varphi} : B_n \rightarrow \mathbb{R}$ a homogeneous quasi-morphism, and $\tilde{\Phi}$ is the corresponding homogeneous quasi-morphism on $\mathcal{D}$. Then*

$$\tilde{\Phi}(h_1) = \int_{T_H} \tilde{h}' d\mu_{\tilde{\varphi}}, \quad (25)$$

where h_1 is the time-one Hamiltonian flow generated by H .

Sketch of the proof. Suppose that the corresponding Reeb graph T_H has l edges.

Take an n -tuple $\bar{x} = (x_1, \dots, x_n) \in X_n$ so that each x_i is not a critical point of H , and $H(x_i) \neq H(x_j)$ for each x_i, x_j such that the corresponding points in T_H lie on the same edge. For the simplicity we will assume that there is a sequence of natural numbers $\{n_i\}_{i=1}^l$ such that $\sum_{i=1}^l n_i = n$, and the projections of the points $x_1, \dots, x_{n_1}$ on T_H lie on the edge e_1 , the projections of the points $x_{n_1+1}, \dots, x_{n_1+n_2}$ on T_H lie on the edge e_2 and so on.

Let h_t be a Hamiltonian flow generated by H . Take $p \in \mathbb{N}$. Then

$$\gamma(h_1^p; \bar{x}) = \alpha_{1,p} \gamma_{1,p}(h_1; \bar{x}) \cdot \dots \cdot \gamma_{l,p}(h_1; \bar{x}) \alpha_{2,p}, \quad (26)$$

for some $\gamma_{i,p}(h_1; \bar{x}) \in P_n$, which pairwise commute with each other, and for some $\alpha_{1,p}$ and $\alpha_{2,p}$. The braids $\alpha_{1,p} \in B_n$ and $\alpha_{2,p} \in B_n$ depend on $\bar{x} \in X_n$, $p \in \mathbb{N}$ and H . For all $n \in \mathbb{N}$ there exists a constant $M(n) > 0$, which depends only on n , such that for each $\bar{x} \in X_n$ as above and for each $p \in \mathbb{N}$ this constant satisfies the following inequations: $|\tilde{\varphi}(\alpha_{1,p})| < M(n)$, $|\tilde{\varphi}(\alpha_{2,p})| < M(n)$. Each $\gamma_{i,p}(h_1; \bar{x})$ depends on the integer p , the function H , the points

$$x_{n_1+n_2+\dots+n_{i-1}+1}, \dots, x_{n_1+n_2+\dots+n_i},$$

and on the position of the points whose projections on T_H do not lie on the edge e_i .

For $\beta \in P_k$ we define $\beta_{k,m} \in P_{k+m}$ to be an image of β under the standard inclusion of P_k into P_{m+k} . It means that $\beta_{k,m} = \beta \sqcup Id_m$, where the identity braid $Id_m \in P_m$ is placed on the right of the braid $\beta \in P_k$. We denote

$$\beta(i, p) = \gamma_{i,p}(h_1; x_{n_1+n_2+\dots+n_{i-1}+1}, \dots, x_{n_1+n_2+\dots+n_i}).$$

It is easy to see that each $\gamma_{i,p}(h_1; \bar{x})$ is conjugate to the braid $\beta(i, p)_{n_i, n-n_i}$. Thus the conjugacy class of each $\gamma_{i,p}(h_1; \bar{x})$ depends only on the integer p , the function H and the points $x_{n_1+n_2+\dots+n_{i-1}+1}, \dots, x_{n_1+n_2+\dots+n_i}$.

Note that any homogeneous quasi-morphism restricts to a homomorphism on an abelian subgroup, and is constant on any conjugacy class. Thus

$$\lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\gamma(h_1^p; \bar{x})) = \sum_{i=1}^l \lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\gamma_{i,p}(h_1; \bar{x})) \quad (27)$$

and

$$\begin{aligned} \tilde{\Phi}(h_1) &= \int_{X_n} \lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\gamma(h_1^p; \bar{x})) d\bar{x} = \sum_{i=1}^l \int_{X_n} \lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\gamma_{i,p}(h_1; \bar{x})) d\bar{x} = \\ &= \sum_{i=1}^l \sum_{\substack{n_1, \dots, n_l \\ \sum_{j=1}^l n_j = n}} \int_{X_n} \lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\beta(i, p)_{n_i, n-n_i}) d\bar{x}. \end{aligned} \quad (28)$$

Note that

$$\int_{X_n} \lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\beta(i, p)_{n_i, n-n_i}) d\bar{x} = \text{vol} \left(\prod_{\substack{j=1 \\ j \neq i}}^l \prod_{k=1}^{n_j} A_j \right) \int_{X_{A_i, n_i}} \lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\beta(i, p)_{n_i, n-n_i}) d\bar{x}_{n_i}, \quad (29)$$

where $d\bar{x}_{n_i} = dx_{n_1+n_2+\dots+n_{i-1}+1} \cdot \dots \cdot dx_{n_1+n_2+\dots+n_{i-1}+n_i}$, A_j is an annulus which corresponds to an edge e_j for all $0 \leq j \leq l$, and X_{A_i, n_i} is the space of ordered n_i -tuples of different points in A_i .

Denote the last integral in Equation (29) by $\mathfrak{I}_i$. Thus equations (28) and (29) tell us that in order to compute $\tilde{\Phi}(h_1)$ it is enough to know the value $\mathfrak{I}_i$ for each $1 \leq i \leq l$. We will compute $\mathfrak{I}_i$ by applying action-angle coordinates on the domain A_i .

From now on we will discuss only the model case, where the function $H|_{\mathbb{D}^2 \setminus \overline{U}}$ has 2 maximum points and 1 saddle point. The level curves of H are presented in Figure 6. Let us cut the disc into 4 pieces, by cutting it along its critical level curves, which passes through the critical points, and by cutting it along a curve c and $\partial \mathbb{D}^2$. We get 4 open annuli. Denote the 3 open annuli, where H is not identically zero by A_1, A_2 ,




Figure 6: Level curves

and A_3 . The open annulus A_3 is such that the curve c is a connected component of ∂A_3 . We denote by A_0 the other open annulus, see Figure 6. Set $R_1 = \frac{\text{area}(A_1)}{2\pi}$ and $R_2 = \frac{\text{area}(A_2)}{2\pi}$. Let (I_i, θ_i) be angle-action coordinates on A_i for $1 \leq i \leq 3$, and let (I_0, θ_0) be the coordinates on A_0 , where I_0 is the area coordinate divided by 2π and θ_0 is the angle coordinate. Set $w_i^j = \frac{\partial h_i}{\partial I_i^j}(I_i^j)$, where $1 \leq i \leq 3$ and $1 \leq j \leq n$, and note that $w_i^j \leq 0$ for all $1 \leq i \leq 3$ and $1 \leq j \leq n$, because H has two maximum points.

Let $\eta_{i,k,n}^+$ be a pure braid in P_n , in which the $i+k$ -th strand makes one twist in the positive direction around previous $i-1$ strands, and let $\eta_{i,k,n}^-$ be a pure braid in P_n , in which the $i+k$ -th strand makes one twist in the negative direction around previous $i-1$ strands (see Figure 7). If $k=0$, then instead of $\eta_{i,0,n}$ we will write $\eta_{i,n}$.

For each $0 \leq k_1 \leq n$, $0 \leq k_2 \leq n$, $0 \leq k_3 \leq n$, $0 \leq k_1 + k_2 \leq k_3 \leq n$ the set:

$$A_{k_1, k_2, k_3, n} = \{\eta_{i,n}^+, \eta_{i,n}^-, \eta_{j,k_1,n}^+, \eta_{j,k_1,n}^-, \eta_{k,n}^+, \eta_{k,n}^- | 2 \leq i \leq k_1, 2 \leq j \leq k_2, k_1 + k_2 + 1 \leq k \leq k_3\}$$

generates an abelian subgroup of P_n . It is easy to see that for each $2 \leq j \leq k_2$ the braids $\eta_{j,k_1,n}$ and $\eta_{j,n}$ are conjugate. Thus $\tilde{\varphi}(\eta_{j,k_1,n}) = \tilde{\varphi}(\eta_{j,n})$.

Take the points $(I_1^1, \theta_1^1), \dots, (I_1^{k_1}, \theta_1^{k_1}) \in A_1$ such that $I_1^1 < \dots < I_1^{k_1}$. Take the points $(I_2^1, \theta_2^1), \dots, (I_2^{k_2}, \theta_2^{k_2}) \in A_2$ such that $I_2^1 < \dots < I_2^{k_2}$. Take the points $(I_3^{k_1+k_2+1}, \theta_3^{k_1+k_2+1}), \dots, (I_3^{k_3}, \theta_3^{k_3}) \in A_3$ such that $I_3^{k_1+k_2+1} < \dots < I_3^{k_3}$.




Figure 7: The braid $\eta_{i,k,n}^+$

Take the points $(I_0^{k_3+1}, \theta_0^{k_3+1}), \dots, (I_0^n, \theta_0^n) \in A_0$ such that $I_0^{k_3+1} < \dots < I_0^n$. We denote

$$\bar{I} = (I_1^1, \dots, I_1^{k_1}, I_2^1, \dots, I_2^{k_2}, I_3^{k_1+k_2+1}, \dots, I_3^{k_3}, I_0^{k_3+1}, \dots, I_0^n),$$

and

$$\bar{\theta} = (\theta_1^1, \dots, \theta_1^{k_1}, \theta_2^1, \dots, \theta_2^{k_2}, \theta_3^{k_1+k_2+1}, \dots, \theta_3^{k_3}, \theta_0^{k_3+1}, \dots, \theta_0^n).$$

For each $p \in \mathbb{N}$ the pure braid

$$\begin{aligned} \gamma(h_1^p; \bar{I}, \bar{\theta}) &:= \gamma(h_1^p; (I_1^1, \theta_1^1), \dots, (I_1^{k_1}, \theta_1^{k_1}), (I_2^1, \theta_2^1), \dots, (I_2^{k_2}, \theta_2^{k_2}), \\ &\quad (I_3^{k_1+k_2+1}, \theta_3^{k_1+k_2+1}), \dots, (I_3^{k_3}, \theta_3^{k_3}), (I_0^{k_3+1}, \theta_0^{k_3+1}), \dots, (I_0^n, \theta_0^n)) \end{aligned}$$

reads as:

$$\alpha_{1,p} \eta_{2,n}^{-[-pw_1^2]} \dots \eta_{k_1,n}^{-[-pw_1^{k_1}]} \eta_{2,k_1,n}^{-[-pw_2^2]} \dots \eta_{k_2,k_1,n}^{-[-pw_2^{k_2}]} \eta_{k_1+k_2+1,n}^{-[-pw_3^{k_1+k_2+1}]} \dots \eta_{k_3,n}^{-[-pw_3^{k_3}]} \alpha_{2,p},$$

where $[\cdot]$ stands for the integer part.

Using the fact that $\eta_{i,k,n}^- = (\eta_{i,k,n}^+)^{-1}$, $\tilde{\varphi}(\eta_{i,k,n}) = \tilde{\varphi}(\eta_{i,n})$, and that $\tilde{\varphi}$ is a homogeneous quasi-morphism, we get:

$$\lim_{p \rightarrow \infty} \frac{1}{p} \tilde{\varphi}(\gamma(h_1^p; \bar{I}, \bar{\theta})) = \sum_{i=2}^{k_1} \tilde{\varphi}(\eta_{i,n}) w_1^i + \sum_{i=2}^{k_2} \tilde{\varphi}(\eta_{i,n}) w_2^i + \sum_{i=k_1+k_2+1}^{k_3} \tilde{\varphi}(\eta_{i,n}) w_3^i.$$

Denote

$$\tilde{\varphi}_{k_1,k_2,k_3} = \sum_{i=2}^{k_1} \tilde{\varphi}(\eta_{i,n}) w_1^i + \sum_{i=2}^{k_2} \tilde{\varphi}(\eta_{i,n}) w_2^i + \sum_{i=k_1+k_2+1}^{k_3} \tilde{\varphi}(\eta_{i,n}) w_3^i,$$

and

$$dI = dI_1^1 \dots dI_1^{k_1} dI_2^1 \dots dI_2^{k_2} dI_3^{k_1+k_2+1} \dots dI_3^{k_3} dI_0^{k_3+1} \dots dI_0^n.$$

Thus

$$\tilde{\Phi}(h_1) = n!(2\pi)^n \sum_{k_1=0}^n \sum_{k_2=0}^{n-k_1} \sum_{k_3=k_1+k_2+1}^n \int \cdots \int_{D_{k_1,k_2,k_3}} \tilde{\varphi}_{k_1,k_2,k_3} dI,$$

where $D_{k_1,k_2,k_3} \subset [0, \frac{1}{2}]^{\times n}$ is the domain which is determined by the following inequalities: $0 < I_1^1 < \dots < I_1^{k_1} < R_1$, $0 < I_2^1 < \dots < I_2^{k_2} < R_2$, $R_1 + R_2 < I_3^{k_1+k_2+1} < \dots < I_3^{k_3} < \frac{1}{2} - \varepsilon$ and $\frac{1}{2} - \varepsilon < I_0^{k_3+1} < \dots < I_0^n < \frac{1}{2}$.

We had to multiply by $(2\pi)^n$, because there are n angle coordinates. In order to count all the ordered n -tuples of points we had to multiply by $n!$. After a simple computation (see [7]) we get

$$\tilde{\Phi}(h_1) = \int_{T_H} \hbar' d\mu_{\tilde{\varphi}}. \quad (30)$$

□

Proof of Theorem 4.8. The theorem of Banyaga [3] states that $\ker(\mathcal{C})$ is a simple group, where $\mathcal{C} : \mathcal{D} \rightarrow \mathbb{R}$ is a Calabi homomorphism (for various definitions of this homomorphism see [8], [11] and [20]). Thus there exists a homomorphism $v : \mathbb{R} \rightarrow \mathbb{R}$ such that for all $g \in \mathcal{D}$

$$v(\mathcal{C}(g)) = \widetilde{Ras}_n(g). \quad (31)$$

Restricting $\widetilde{Ras}_n$ and $\mathcal{C}$ on a 1-parametric subgroup of $\mathcal{D}$, on which they do not vanish, one easily sees that v is continuous. It means that

$$\mathcal{K}_n \cdot \mathcal{C}(g) = \widetilde{Ras}_n(g), \quad (32)$$

where $\mathcal{K}_n$ is some constant in $\mathbb{R}$ which depends only on n .

Take $H : \mathbb{D}^2 \rightarrow \mathbb{R}$, such that H equals zero near the boundary, and such that H has two maximum points and one saddle point. The level curves of H are shown in Figure 6. Denote by g the time-one Hamiltonian flow generated by H . An explicit calculation of $\widetilde{Ras}_n$ using Theorem 5.4 and $\mathcal{C}$ on g shows that the homomorphism v has the form

$$v(s) = 2n(n-1)\pi^{n-1} \cdot s.$$

□

Remark 5.5. The formula (25) fits in with Theorem 4.8 when $\tilde{\varphi} = lk$ is a canonical homomorphism

$$lk : B_n \rightarrow \mathbb{Z},$$

such that $lk(\sigma_i) = 1$, $lk(\sigma_i^{-1}) = -1$ and σ_i is a standard generator of B_n .

Remark 5.6. For each $n > 1$ the homogeneous quasi-morphism $\widetilde{\Phi}_n$, defined by a homogeneous quasi-morphism $\widetilde{\varphi} : P_n \rightarrow \mathbb{R}$, is not continuous in C^0 -topology. The case $n = 2$ was proved by Gambaudo and Ghys in [11]. The proof in the case $n > 2$ is almost identical to their proof.

Recall that $\text{sign}_n : B_n \rightarrow \mathbb{R}$ is a quasi-morphism for all $n \geq 2$. The induced family of homogeneous quasi-morphisms was denoted by $\widetilde{\text{Sign}}_{n, \mathbb{D}^2}$. In [12] Gambaudo and Ghys proved that all of them are non-trivial and linearly independent. In the case where the Reeb graph of H is simple they showed that

$$\widetilde{\text{Sign}}_{n, \mathbb{D}^2}(h_1) = \int_{T_H} \hbar' d\mu_{\widetilde{\varphi}_n}.$$

Proposition 5.7. *For each natural number n we have*

$$\widetilde{\text{Sign}}_{n, \mathbb{D}^2}(h_1) = n\pi^n \int_{T_H} ((1 - 4I)^{n-1} - 4(n-1)I - 1) \hbar'(I) dI, \quad (33)$$

where h_1 is the time-one Hamiltonian flow defined by a Morse type Hamiltonian H .

Proof. Recall that $\eta_{i,n}$ is a pure braid in P_n , in which the i -th strand makes one twist in the positive direction around previous $i - 1$ strands. Using the fact (see page 148 in [22]) that

$$\widetilde{\text{sign}}_n(\eta_{i,n}) = \begin{cases} -i, & \text{if } i \text{ is even,} \\ 1-i, & \text{if } i \text{ is odd} \end{cases}$$

it is easy to prove the following formula:

$$\begin{aligned} \sum_{i=2}^n \widetilde{\text{sign}}_n(\eta_{i,n}) i \binom{n}{i} x^{i-1} \left(\frac{1}{2} - x\right)^{n-i} = \\ \frac{n}{2} \left(\left(\frac{1}{2} - 2x\right)^{n-1} - 4(n-1)x \left(\frac{1}{2}\right)^{n-1} - \left(\frac{1}{2}\right)^{n-1} \right). \end{aligned} \quad (34)$$

Now the proof follows easily from Theorem 5.4 and (34). $\square$

Proof of Proposition 4.2. Claim 1. Take a braid $\alpha_n \in \underline{B}_n$ for $n > 2$, where $\alpha_n = \sigma_1 \cdots \sigma_{n-1}$. Then it follows from [[22], page 148] that $\widetilde{\text{sign}}_n(\alpha_n) = -\frac{n}{2}$ if n is even, and $\widetilde{\text{sign}}_n(\alpha_n) = -\frac{n^2-1}{2n}$ if n is odd. Thus if n is even then:

$$-\frac{n}{2} = \widetilde{\text{sign}}_n(\alpha_n) \neq \widetilde{\text{sign}}_n(\alpha_{n-1}) + \widetilde{\text{sign}}_n(\sigma_{n-1}) = -\frac{(n-1)^2-1}{2n-2} - 1,$$

and if n is odd then:

$$-\frac{n^2-1}{2n} = \widetilde{\text{sign}}_n(\alpha_n) \neq \widetilde{\text{sign}}_n(\alpha_{n-1}) + \widetilde{\text{sign}}_n(\sigma_{n-1}) = -\frac{n-1}{2} - 1.$$

Claim 2. Suppose that $\widetilde{\text{sign}}_n : P_n \rightarrow \mathbb{R}$ is a homomorphism. Then $\widetilde{\text{Sign}}_{n, \mathbb{D}^2} : \mathcal{D} \rightarrow \mathbb{R}$ is a homomorphism. Let $\mathcal{C} : \mathcal{D} \rightarrow \mathbb{R}$ be the Calabi homomorphism. Using the theorem of Banyaga [3], which states that $\ker(\mathcal{C})$ is a simple group, we get that there exists a homomorphism $\kappa : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\widetilde{\text{Sign}}_{n, \mathbb{D}^2} = \kappa \mathcal{C}.$$

Restricting $\widetilde{\text{Sign}}_{n, \mathbb{D}^2}$ and $\mathcal{C}$ on a 1-parametric subgroup of $\mathcal{D}$, on which they do not vanish, one easily sees that κ is continuous. Hence $\kappa(y) = K_n y$ for all $y \in \mathbb{R}$, and

$$\widetilde{\text{Sign}}_{n, \mathbb{D}^2} = K_n \mathcal{C},$$

where K_n is some real constant which depends only on n . But using Proposition 5.7, we see that this is impossible for $n > 2$. $\square$

Proof of Theorem 4.3. Step 1. Suppose that g is generated by a Morse type Hamiltonian $H \in \mathcal{H}_0$. Then using (33) we get after a simple computation that

$$\lim_{n \rightarrow \infty} \frac{\widetilde{\text{Sign}}_{n, \mathbb{D}^2}(g)}{\pi^{n-1} n(n-1)} = -\mathcal{C}(g).$$

Step 2. Suppose that g is generated by any $H \in \mathcal{H}_0$. Then there exists a Morse type function $F \in \mathcal{H}_0$, which is C^1 -close to H (see Theorem 2.7 in [21]). Now the statement follows from Theorem 5.11. $\square$

The goal of the remaining part of this section is to show that if $H_n \xrightarrow[n \rightarrow \infty]{} H$ in C^1 -topology, then $\lim_{n \rightarrow \infty} \widetilde{\Phi}(h_{1,n}) = \widetilde{\Phi}(h_1)$. Here $H_n \in \mathcal{H}_0$, $H \in \mathcal{H}_0$, and $h_{1,n}$, h_1 are Hamiltonian time-one flows generated by H_n and H respectively.

Let $g \in \mathcal{D}$ and $\psi_t : [0, 1] \rightarrow \mathcal{D}$ be such that $\psi_0 = Id$ and $\psi_1 = g$. We denote

$$\mathcal{G}(\{\psi_t\}) = \int_{\mathbb{D}^2 \times \mathbb{D}^2} \frac{1}{2\pi} \int_0^1 \left\| \frac{\partial}{\partial t} \left(\frac{\psi_t(x) - \psi_t(y)}{\|\psi_t(x) - \psi_t(y)\|} \right) \right\| dt dx dy.$$

Lemma 1 in [14] shows that $\mathcal{G}(\{\psi_t\})$ is well defined. Denote by $\mathcal{G}(g) = \inf \mathcal{G}(\{\psi_t\})$, where the infimum is taken over all isotopies in $\mathcal{D}$ joining the identity with g . Lemma 2 in [14] states that for all g and h in $\mathcal{D}$

$$\mathcal{G}(gh) \leq \mathcal{G}(g) + \mathcal{G}(h).$$

Thus the following limit exists:

$$\mathcal{L}(g) = \lim_{n \rightarrow \infty} \frac{\mathcal{G}(g^n)}{n}.$$

Proposition 5.8. *Let $g \in \mathcal{D}$, and let $\varphi : B_n \rightarrow \mathbb{R}$ be a quasi-morphism. Then:*

$$|\tilde{\Phi}(g)| \leq K_1 \mathcal{L}(g),$$

where $K_1 > 0$ depends only on n .

Proof. From the proof of Lemma 2.1 we get the following inequalities:

$$|\tilde{\Phi}(g)| \leq \lim_{p \rightarrow \infty} \frac{1}{p} \int_{X_n} |\tilde{\varphi}(\gamma(g^p; \bar{x}))| d\bar{x} \leq 2N \cdot \text{vol}((\mathbb{D}^2)^{n-2}) \sum_{i < j}^n \mathcal{L}(g) \leq K_1 \mathcal{L}(g),$$

where $K_1 = nN(n+1)\text{vol}((\mathbb{D}^2)^{n-2})$. □

Now we will define a metric on $\mathcal{D}$. A tangent vector to $\mathcal{D}$ at a point g is a Hamiltonian vector field X_g (see [9]) whose L^2 -norm is defined by:

$$\|X_g\|_2 = \left(\int_{\mathbb{D}^2} \|X_g(x)\|^2 dx \right)^{\frac{1}{2}},$$

where $\|\cdot\|$ stands for the standard Euclidean norm. For any path ψ_t in $\mathcal{D}$ which connects g and h , the length of this path is given by the formula:

$$l(\{\psi_t\}) = \int_0^1 \left\| \frac{d\psi_t}{dt} \right\|_2 dt.$$

We define by $d_2(g, h) = \inf l(\{\psi_t\})$, where the infimum is taken over all paths in $\mathcal{D}$ joining g with h . One can show that d_2 is a distance function. Thus $(\mathcal{D}, d_2)$ is a metric space. Note that d is right invariant: $d_2(g, h) = d_2(gf, hf)$.

Corollary 5.9. *Let $g \in \mathcal{D}$, and let $\varphi : B_n \rightarrow \mathbb{R}$ be a quasi-morphism. Then:*

$$|\tilde{\Phi}(g)| \leq K_2 d_2(\text{Id}, g),$$

where $K_2 > 0$ depends only on n .

Proof. Theorem 1 in [14] says that for any $g \in \mathcal{D}$ there exists some universal constant $K > 0$, such that

$$\mathcal{L}(g) \leq K d_2(\text{Id}, g).$$

Now take $K_2 = K \cdot K_1$. The statement follows from Proposition 5.8. □

Lemma 5.10. *Let $F \in \mathcal{H}_0$. Then for any $\varepsilon > 0$ and $p \in \mathbb{N}$ there exists $\delta_p > 0$, such that if $H \in \mathcal{H}_0$ is δ_p -close to F in C^1 -topology, then*

$$d_2(f_1^p, h_1^p) < \varepsilon,$$

where f_t and h_t are Hamiltonian flows generated by F and H .

Proof. For the convenience we normalize the area of $\mathbb{D}^2$ to be 1. First, we will show that for all $p \in \mathbb{N}$ there exists $\delta_{1,p} > 0$ such that

$$\max_{x \in \mathbb{D}^2} \|\nabla F(x) - \nabla H(x)\| < \delta_{1,p} \Rightarrow d_2(f_1^p, h_1^p) < \varepsilon.$$

Note that

$$d_2(f_1^p, h_1^p) = d_2(\text{Id}, f_1^p h_1^{-p}) \leq l(\{f_t^p h_t^{-p}\}).$$

It follows from Proposition 1.4.D in [24] that

$$\frac{\partial(f_t^p h_t^{-p})}{\partial t}(x) = p \cdot (\text{sgrad}(F) - \text{sgrad}(H f_t^{-p}))_{(f_t^p h_t^{-p}(x))},$$

where sgrad is the symplectic gradient. Thus

$$\begin{aligned} \left\| \frac{\partial(f_t^p h_t^{-p})}{\partial t}(x) \right\| &= p \cdot \left\| (\text{sgrad}(F) - \text{sgrad}(H f_t^{-p}))_{(f_t^p h_t^{-p}(x))} \right\| \\ &= p \cdot \left\| (\nabla F - \nabla(H f_t^{-p}))_{(f_t^p h_t^{-p}(x))} \right\|, \end{aligned}$$

Denote by $M_{Df_t} = \max_{x \in \mathbb{D}^2, t \in [0,1]} \|(Df_t^{-1})_{(x)}\|$. Note that f_t is an autonomous Hamiltonian flow. Thus $F f_t(x) = F(x)$ for all $x \in \mathbb{D}^2$ and $t \in \mathbb{R}$. It follows that $\forall x \in \mathbb{D}^2$ and $\forall p \in \mathbb{Z}$

$$\nabla F_{(h_t^{-p}(x))} (Df_t^{-p})_{(f_t^p h_t^{-p}(x))} = \nabla F_{(f_t^p h_t^{-p}(x))}.$$

We get the following inequality:

$$d_2(f_1^p, h_1^p) \leq p \int_0^1 \left(\int_{\mathbb{D}^2} \|(Df_t^{-p})_{(f_t^p h_t^{-p}(x))}\|^2 \|\nabla F_{(h_t^{-p}(x))} - \nabla H_{(h_t^{-p}(x))}\|^2 \right)^{\frac{1}{2}} dt.$$

Take $\delta_{1,p} = \frac{\varepsilon}{p} (M_{Df_t})^p$. We get the following inequality

$$d_2(f_1^p, h_1^p) \leq p (M_{Df_t})^p \max_{x \in \mathbb{D}^2} \|\nabla F(x) - \nabla H(x)\| < \varepsilon.$$

Denote by $\delta_p = \frac{\delta_{1,p}}{2}$. So we have shown that if H is δ_p -close to F in C^1 -topology then $d_2(f_1^p, h_1^p) < \varepsilon$. $\square$

Theorem 5.11. *Let $F \in \mathcal{H}_0$. Then for any ε , such that $1 > \varepsilon > 0$, there exists $\delta > 0$ such that if $H \in \mathcal{H}_0$ is δ -close to F in C^1 -topology then:*

$$\left| \tilde{\Phi}(f_1) - \tilde{\Phi}(h_1) \right| \leq \varepsilon,$$

where f_t and h_t are Hamiltonian flows generated by F and H .

Proof. Let $0 < \varepsilon < 1$, let $D_{\tilde{\Phi}}$ be the defect of the homogeneous quasi-morphism

$$\tilde{\Phi} : \mathcal{D} \rightarrow \mathbb{R},$$

and let K_2 be the constant which was defined in Corollary 5.9. Take $p \in \mathbb{N}$ such that $\frac{D_{\tilde{\Phi}} + K_2}{p} < \varepsilon$. It follows from Lemma 5.10 that there exists $\delta_p > 0$, such that if H is δ_p -close to F in C^1 -topology, then $d_2(f_1^p, h_1^p) < 1$. Thus we get the following inequality:

$$\left| \tilde{\Phi}(f_1) - \tilde{\Phi}(h_1) \right| = \frac{1}{p} \left| \tilde{\Phi}(f_1^p) - \tilde{\Phi}(h_1^p) \right| \leq \frac{D_{\tilde{\Phi}} + \left| \tilde{\Phi}(f_1^p h_1^{-p}) \right|}{p}.$$

It follows from Corollary 5.9 that:

$$\left| \tilde{\Phi}(f_1^p h_1^{-p}) \right| \leq K_2 d_2(\text{Id}, f_1^p h_1^{-p}) = K_2 d_2(f_1^p, h_1^p) < K_2.$$

Thus

$$\left| \tilde{\Phi}(f_1) - \tilde{\Phi}(h_1) \right| < \frac{D_{\tilde{\Phi}} + K_2}{p} < \varepsilon.$$

□

6 Open problems

The following three open problems arise naturally in connection with our results.

Problem 1. Does there exist a knot invariant I , independent from the signatures, τ and the Rasmussen invariant s , such that I satisfies the conditions of Theorem 3.3, and such that the homogenization of the induced quasi-morphism $I : P_n \rightarrow \mathbb{R}$ is non-trivial?

Problem 2. Surprisingly, the proof of the second claim in Proposition 4.2 – dealing with a quasi-morphism on P_n – involves the Calabi homomorphism on $\mathcal{D}$. It would be interesting to find a direct proof of this claim.

Problem 3. The following problem is motivated by a question posed to us by L. Polterovich. Denote by $\mathcal{D}_{\text{aut}} \subset \mathcal{D}$ the set of area-preserving diffeomorphisms generated by autonomous Hamiltonians.

Observe that, by our results in Section 4.4, one can find a linear combination Φ of homogeneous quasi-morphisms $\widetilde{\text{sign}_\omega}$ on B_n so that the corresponding homogeneous quasi-morphism $\tilde{\Phi} : \mathcal{D} \rightarrow \mathbb{R}$ vanishes on $\mathcal{D}_{\text{aut}}$.

On the other hand, using a construction from [12] (completely different from the one described in this paper) one can construct more homogeneous quasi-morphisms $\tilde{\Psi}_\alpha$ on $\mathcal{D}$ that vanish on $\mathcal{D}_{aut}$.

It would be interesting to check whether the quasi-morphism $\tilde{\Phi}$ is non-trivial and, if so, whether it is a linear combination of the quasi-morphisms $\tilde{\Psi}_\alpha$.

So far we have not been able to compute explicitly the value of *any* of the Gambaudo-Ghys quasi-morphisms coming from quasi-morphisms on B_n on *any* area-preserving diffeomorphism that is not generated by a Hamiltonian flow preserving a foliation.

References

- [1] Arnold V. I.: *Mathematical methods of classical mechanics*, translated from the 1974 Russian edition by K. Vogtmann and A. Weinstein, Graduate Texts in Math., Vol. 60, Springer-Verlag, New York and Berlin, 1978.
- [2] Baader S.: *Asymptotic Rasmussen invariant*, arXiv: math.GT/0702335, 2007.
- [3] Banyaga A.: *Sur la structure du groupe des difféomorphismes qui préservent une forme symplectique*, (French) Comment. Math. Helv. 53 (1978), no. 2, 174-227.
- [4] Barge, J., Ghys, E., *Cocycles d'Euler et de Maslov*, Math. Ann. 294:2 (1992), 235-265.
- [5] Beliakova A., Wehrli S.: *Categorification of the colored Jones polynomial and Rasmussen invariant of links*, arXiv: math.GT/0510382, 2006.
- [6] Biran P., Entov M., Polterovich L.: *Calabi quasimorphisms for the symplectic ball*, Communications in Contemporary Mathematics, 6:5 (2004), 793-802.
- [7] Brandenbursky M.: *Ph. D. thesis*, in preparation.
- [8] Calabi E.: *On the group of automorphisms of a symplectic manifold*, Problems in Analysis, symposium in honour of S. Bochner, ed. R.C. Gunning, Princeton Univ. Press (1970), 1-26.
- [9] Ebin, D. G., Marsden, J.: *Groups of diffeomorphisms and the motion of an incompressible fluid*, Ann. of Math. 92, (1970), 102-163.
- [10] Fathi A., Laudenbach F., Poenaru V.: *Travaux de Thurston sur les surfaces*. Astérisque, 66-67. Societe Mathematique de France, Paris, (1979).
- [11] Gambaudo J.M., Ghys E.: *Enlacements asymptotiques*, Topology 36 (1997), no. 6, 1355-1379.

- [12] Gambaudo J.M., Ghys E.: *Commutators and diffeomorphisms of surfaces*, Ergodic Theory Dynam. Systems 24 (2004), no. 5, 1591-1617.
- [13] Gambaudo J.M., Ghys E.: *Braids and signatures*, Bull. Soc. Math. France 133 (2005), no. 4, 541-579.
- [14] Gambaudo J.M., Lagrange M.: *Topological lower bounds on the distance between area preserving diffeomorphisms*, Bol. Soc. Brasil. Mat. 31, (2000), 1-19. CMP 2000:12
- [15] Grigorchuk R., *Some results on bounded cohomology*, Combinatorial and Geometric Group Theory, LMS Lecture Notes Ser. 284, Cambridge University Press, (1994) 111-163.
- [16] Khovanov M.: *A categorification of the Jones polynomial*, Duke Math J., 101 (2000), 359-426.
- [17] Lee E. S.: *An endomorphism of the Khovanov invariant*. arXiv: math.GT/0210213, 2004.
- [18] W. B. Raymond Lickorish: *An Introduction to Knot Theory*. 1997 Springer-Verlag New York, Inc.
- [19] Livingston C.: *Computations of the Ozsvath-Szabo knot concordance invariant*, Geometry and Topology 8 (2004) 735-742.
- [20] McDuff D., Salamon D.: *Introduction to Symplectic Topology*, Oxford Science Publications.
- [21] Milnor, John W.: *Lectures on the h-cobordism theorem*, notes by L. Siebenmann and J. Sondow, Princeton University Press, Princeton, NJ, (1965).
- [22] Murasugi K.: *Knot theory and its applications*, Translated from the 1993 Japanese original by Bohdan Kurpita. Birkhäuser Boston, Inc., Boston, (1996).
- [23] Ozsvath P. S. and Szabo Z.: *Knot Floer homology and the four-ball genus*, Geometry and Topology 7 (2003) 615-639
- [24] Polterovich L.: *The geometry of the group of symplectic diffeomorphisms*, Lectures in Mathematics ETH Zurich, Birkhauser, Basel, 2001. xii+132 pages.
- [25] Rasmussen J. A.: *Khovanov homology and the slice genus*, arXiv: math.GT/0402131, 2004.
- [26] Shumakovitch A.: *Rasmussen invariant, slice-Bennequin inequality, and sliceness of knots*, arXiv: math.GT/0411643, 2004.

Michael Brandenbursky
Department of Mathematics
Technion - Israel Institute of Technology
Haifa 32000, Israel
E-mail address: `mbranden@tx.technion.ac.il`